

Comment on “Supercurrent Noise in Quantum Point Contacts”

In a recent letter, Averin and Imam [1] have analyzed the current fluctuations in a superconducting quantum point contact (SQPC). Their analysis shows that the current noise can be extremely large due to the interplay between quasi-particle scattering and superconducting coherence. This conclusion is in agreement with the results given by us in a recent publication [4].

There are, however, certain limitations in the analysis given in Ref. [1] which, as we discuss in this comment, can lead to the disappearance of an important feature in the low frequency noise-spectrum.

The main results of Ref. [1] are obtained for the specific case of a perfectly transmitting contact in which certain simplifying relations hold, allowing Averin and Imam to give an appealing and simple real time interpretation of the current fluctuations in a SQPC. However, for the analysis of real experimental situations involving a few quantum channels, like the ones in Refs. [2,3], a discussion of the *nearly* perfect transmission or quasi-ballistic case is desirable. In fact, it turns out that even a small departure from perfect transmission leads to the appearance of a relevant new structure in the noise spectral density.

In Ref. [4] we have calculated the supercurrent noise spectrum $S(\omega)$ of a SQPC starting from a microscopic model with arbitrary transmission. We found that for the general case $S(\omega)$ exhibits a pronounced resonant peak at $\omega = 2\omega_S$, where ω_S is the subgap state energy, in *addition* to the zero-frequency peak shown in Fig. 1 of Ref. [1] for perfect transmission. The overall two-peaked structure of $S(\omega)$ is illustrated in Fig. 1 of Ref. [4]. The relative weight of the peak at $2\omega_S$ with respect to the one at zero-frequency can be calculated from the analytical expressions given in Ref. [4], yielding

$$S(2\omega_S)/S(0) = \frac{1}{2}(1 - \alpha) \tan^2 \frac{\phi}{2} \cosh \left[\frac{\omega_S}{k_B T} \right] \quad (1)$$

where α is the contact normal transmission and $\omega_S = \Delta \sqrt{1 - \alpha} \sin^2 \phi / 2$.

From this expression it is clear that the peak at $2\omega_S$ can be the dominant feature for low temperatures even when the departure from perfect transmission is very small. Moreover, when $1 - \alpha$ is finite, there are two limits where the only remaining feature is the peak at $2\omega_S$, namely when $\phi \rightarrow \pi$ at finite temperature and when $k_B T \rightarrow 0$.

Finally, let us comment that in Ref. [1] an heuristic argument is given before Eq. (13) for obtaining the extension of $S(\omega)$ to the non-perfect transmission case. Although the neglect of the resonant structure at $2\omega_S$ limits its frequency-range of validity, this argument actually leads to the correct answer for the zero-frequency noise

at any contact transmission, as has been rigorously obtained by the present authors, both by a direct calculation of $S(0)$ [4] and by relating this quantity to the phase-dependent linear conductance via the fluctuation-dissipation theorem [5].

A. Martín-Rodero¹, A. Levy Yeyati¹, F.J. García-Vidal^{1,2} and J.C. Cuevas¹

¹ Departamento de Física Teórica de la Materia Condensada C-V Facultad de Ciencias
Universidad Autónoma de Madrid
E-28049 Madrid, Spain

² Condensed Matter Theory Group
The Blackett Laboratory
Imperial College
London SW7 2BZ, United Kingdom

PACS numbers: 74.50.+r, 73.20.Dx, 74.80.Fp

-
- [1] D. Averin and H.T. Imam, Phys. Rev. Lett. **76**, 3814 (1996).
 - [2] H. Takayanagi, T. Akazaki and J. Nitta, Phys. Rev. Lett. **75**, 3533 (1995).
 - [3] M.C. Koons, L. Feenstra, B.J. Vleeming, A.N. Omelyanchouk and R. de Bruyn Outober, Physica B **218**, 145 (1996).
 - [4] A. Martín-Rodero, A. Levy Yeyati and F.J. García-Vidal, Phys. Rev. B **53**, R8891 (1996).
 - [5] The relation between the phase-dependent linear conductance and $S(0)$ was pointed out in A. Levy Yeyati, A. Martín-Rodero and J.C. Cuevas, Report No. cond-mat 9505102. The detailed calculation of the linear conductance can be found in A. Martín-Rodero, A. Levy Yeyati and J.C. Cuevas, Physica B **218**, 126 (1996); A. Levy Yeyati, A. Martín-Rodero and J.C. Cuevas, J. Phys.: Condens. Matter **8**, 449 (1996).